



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$\text{Then } 17^3 \times 73^3 = \begin{vmatrix} 493 & 340 \\ -340 & 493 \end{vmatrix} \times \begin{vmatrix} 5329 & 0 \\ 0 & 5329 \end{vmatrix} = (493^2 + 340^2)(5329)^2 \\ = (73)^2(493)^2 + (73)^2(340)^2, \text{ which is one set of answers.}$$

Also solved by *R. J. Adcock, and M. A. Gruber.*

19. Proposed by ARTEMAS MARTIN, LL. D., U. S. Coast and Geodetic Survey Office, Washington, D. C.

Find three positive integer numbers whose sum is a cube, and, also, the sum of any two diminished by the third a cube.

I. Solution by M. A. GRUBER, A. M., War Department, Washington, D. C.

Let x , y , and z = the three positive integers.

$$\begin{aligned} \text{Then } x+y+z &= a^3 \\ x+y-z &= b^3 \\ x+z-y &= c^3 \\ y+z-x &= d^3 \end{aligned}$$

$$\text{Whence } x+y+z = b^3 + c^3 + d^3 = a^3;$$

$$x = \frac{b^3 + c^3}{2}; y = \frac{b^3 + d^3}{2}; z = \frac{c^3 + d^3}{2}.$$

This is a problem in which the sum of three cubes = a cube. Take $3^3 + 4^3 + 5^3 = 6^3$. But as the numbers are to be integers, we multiply by 2,

and obtain $6^3 + 8^3 + 10^3 = 12^3$. $\therefore x = \frac{6^3 + 8^3}{2} = 364$; $y = \frac{1}{2}(6^3 + 10^3) = 608$; and

$z = \frac{1}{2}(8^3 + 10^3) = 756$. The number of answers is infinite.

II. Solution by R. J. ADCOCK, Larchland, Warren County, Illinois.

Let x , y , z , be the three numbers; then $x+y+z=u^3$, $x+y-z=r^3$, $x+z-y=s^3$, $z+y-x=t^3$, by conditions.

$$\text{Wherefore } x+y+t=u^3=v^3+r^3+s^3, \quad x=\frac{r^3+r^3}{2}, \quad y=\frac{1}{2}(r^3+s^3),$$

$z=\frac{1}{2}(r^3+s^3)$. The most general equation yet obtained by me for the sum of three cubes = a cube, is found from,

$[(ax^3+dy^3)x]^3 + [(bx^3+hy^3)y]^3 + [(cx^3-hy^3)y]^3 = [(ax^3+gy^3)x]^3$, by expanding, equating coefficients of similar terms with respect to x and y , eliminating d , g , and h , giving the identical equation,

$$[9a^3bx^3y + (b^2-bc+c^2)^2y^4]^3 + [9a^3cx^3y - (b^2-bc+c^2)^2y^4] \\ + [9a^4x^4 - 3(b^2-bc+c^2)caxy^3]^3 = [9a^4x^3 + 3ab(b^2-bc+c^2)xy^3]^3.$$

By numbers for the letters in the above, some of the resulting

equations are $3^3 + 4^3 + 5^3 = 6^3$, $1^3 + 6^3 + 8^3 = 9^3$, $3^3 + 10^3 + 18^3 = 19^3$,

$$7^3 + 14^3 + 17^3 = 20^3, 4^3 + 17^3 + 22^3 = 25^3, 11^3 + 15^3 + 27^3 = 29^3.$$

Then the three positive integer numbers are $x = \frac{11^3 + 15^3}{2}$, $y = \frac{11^3 + 27^3}{2}$,

$z = \frac{15^3 + 27^3}{2}$. Also x, y, z may be found from any equation, including the algebraic sum, for the sum of three cubes = a cube, by first multiplying each cube by 2^3 .

Also solved by *O. W. Anthony, H. W. Draughon, C. D. Schmitt, and G. B. M. Zerr.*

PROBLEMS.

27. Proposed by J. W. NICHOLSON, LL. D., President and Professor of Mathematics, Louisiana State University and A. and M. College, Baton Rouge, Louisiana.

Required a formula for finding five integers the sum of whose cubes is a cube.

28. Proposed by DAVID E. SMITH, Ph. D., Professor of Mathematics, Michigan State Normal School, Ypsilanti, Michigan.

Decompose the product 97.673.257 into the sum of two fourth powers.

AVERAGE AND PROBABILITY.

Conducted by B. F. FINKEL, Kidder, Missouri. All Contributions to this department should be sent to him.

SOLUTIONS OF PROBLEMS.

13. Proposed by I. L. BEVERAGE, Monterey, Virginia.

Find the mean values of the roots of the quadratic $x^2 - ax + b = 0$, the roots being known to be real, but b being unknown and positive.

Solution by P. S. BERG, Apple Creek, Ohio, and JOHN DOLMAN, Jr., Counsellor-at-law, Philadelphia, Penn., and J. M. OULAW, A. M., Principal of High School, Monterey, Virginia.

Solving the given equation, $x = \frac{1}{2}a \pm \sqrt{(\frac{1}{4}a^2 - b)}$.

Therefore, if b be positive and x real, b cannot exceed $\frac{1}{4}a^2$. If β be the smaller of the two roots, its mean value, therefore, is